

MATH227 MATHEMATICAL METHODS FOR NON-PHYSICAL SYSTEMS

JANUARY 2005

1.

$$U(x, y) = xy + 3x + 5y = U_0 \Rightarrow y = \frac{U_0 - 3x}{x + 5} = -3 + \frac{U_0 + 15}{x + 5}.$$

$$\frac{dy}{dx} = -\frac{U_0 + 15}{(x + 5)^2} < 0,$$

$$\frac{d^2y}{dx^2} = 2\frac{U_0 + 15}{(x + 5)^3} > 0,$$

2. Budget constraint touches indifference curve where

$$\frac{\frac{\partial U}{\partial x}}{\frac{\partial U}{\partial y}} = \frac{1}{3}$$

$$\frac{2(x + 1)^2(y + 3)^2}{3(x + 1)^3(y + 3)} = \frac{1}{3}$$

$$2(y + 3) = (x + 1) \Rightarrow x - 2y = 5.$$

Solving with $x + 3y = 10$, we have $x = 7$, $y = 1$. Then $U = U_0 = (7 + 1)^2(1 + 3)^3 = 64 \cdot 64 = 4096$.

3. $c(x, y)$ is minimised where

$$\begin{aligned}\frac{\partial q}{\partial x} &= \frac{\partial c}{\partial x} \\ \frac{\partial q}{\partial y} &= \frac{\partial c}{\partial y} \\ \frac{\frac{4}{5}(x+1)^{-\frac{1}{5}}y^{\frac{1}{5}}}{\frac{1}{5}(x+1)^{\frac{4}{5}}y^{-\frac{4}{5}}} &= \frac{4}{3} \\ 4\frac{y}{(x+1)} &= \frac{4}{3} \Rightarrow y = \frac{1}{3}(x+1) \\ \Rightarrow \left(\frac{1}{3}\right)^{\frac{1}{5}}(x+1) &= 3 \Rightarrow x = 3^{\frac{6}{5}} - 1, \quad y = 3^{\frac{1}{5}} \\ \Rightarrow c = 4x + 3y + 2 &= 4(3^{\frac{6}{5}} - 1) + 3 \cdot 3^{\frac{1}{5}} + 2 = 5.3^{\frac{6}{5}} - 2.\end{aligned}$$

Minimum cost for production level of 3 units is $5.3^{\frac{6}{5}} - 2$ units.

4.

$$C(q) = q^3 - 6q^2 + 14q + 10.$$

(i) Fixed cost $C(0) = 10$. (ii) $MC(q) = C'(q) = 3q^2 - 12q + 14$.
(iii) $AVC(q) = \frac{C(q) - C(0)}{q} = q^2 - 6q + 14$.

Cease production when $p = \min(AVC)$.

$$AVC'(q) = 2q - 6 = 0 \quad \text{when} \quad q = 3 \Rightarrow p = AVC(3) = 5.$$

5.

$$S(p) = 8\frac{p+2}{p+4} \Rightarrow \frac{dS}{dp} = 8\frac{(p+4) - (p+2)}{(p+2)^2} = \frac{16}{(p+2)^2} > 0.$$

Equilibrium where

$$S(p) = D(p) \Rightarrow 8\frac{p+2}{p+4} = \sqrt{48 - 3p}.$$

$p = 4$ is a solution by inspection. Since $S(p)$ is increasing, $D(p)$ is decreasing, it is the only solution. Then amount sold in a week = $D(4) = 6$.

6.

$$C(q) = q^3 - 5q^2 + 9q + 3, \quad D(p) = 12 - p = q \Rightarrow p = 12 - q.$$

Profit given by

$$\begin{aligned} P(q) &= pq - C(q) = (12 - q)q - (q^3 - 5q^2 + 9q + 3) = -q^3 + 4q^2 + 3q - 3 \\ &\Rightarrow P'(q) = -3q^2 + 8q + 3 = -(3q + 1)(q - 3) = 0 \quad \text{for } q = -\frac{1}{3}, \quad 3. \end{aligned}$$

Take the +ve solution $q = 3$. Then $p = 12 - q = 9$.

$$P''(q) = -6q + 8 < 0 \quad \text{for } q = 3.$$

So we have a local maximum. Also

$$P(3) = -3^3 + 4 \cdot 3^2 + 3 \cdot 3 - 3 = 15 > P(0) = -3.$$

So $q = 3$ is a global maximum.

7.

$$\frac{dn}{dt} = -14n + 9n^2 - n^3 = -n(n - 2)(n - 7) = f(n).$$

Equilibrium densities $n = 0$, $n = 2$, $n = 7$.

$f'(0) < 0$, $f(n) \rightarrow -\infty$ as $n \rightarrow \infty$. So graph looks like this:

Equilibria at $n = 0$, $n = 7$ stable; equilibrium at $n = 2$ unstable.

8.

$$\begin{aligned}\frac{dx}{dt} &= x(7 - 3x + y), & \frac{dy}{dt} &= y(7 - x - 2y), \\ 7 - 3x + y &= 7 - x - 2y = 0 \Rightarrow x = 3, & y &= 2, \\ \text{or } y &= 7 - 3x + y = 0 \Rightarrow x = \frac{7}{3}, \\ \text{or } x &= 7 - x - 2y = 0 \Rightarrow y = \frac{7}{2}, \\ \text{or } x &= y = 0.\end{aligned}$$

So the equilibria are $(3, 2)$, $(\frac{7}{3}, 0)$, $(0, \frac{7}{2})$, $(0, 0)$.

9.

$$\mathbf{x} = \mathbf{x}^e + c_1 \mathbf{x}_1 e^{\lambda_1 t} + c_2 \mathbf{x}_2 e^{\lambda_2 t},$$

where $\lambda_{1,2}$ are the e-values, $\mathbf{x}_{1,2}$ are the e-vectors. If λ_1 and λ_2 have opposite signs, then we have a saddle point; the trajectories always move away from \mathbf{x}_e .

10.

$$U(0, N) = 4(N + 1) > U(N, 0) = N + 4$$

so N bars of chocolate preferred to N bags of crisps.

Indifference curves given by

$$y = \frac{U_0 + 2}{x + 4} - 1.$$

Budget constraint touches indifference curve where

$$\frac{\frac{\partial U}{\partial x}}{\frac{\partial U}{\partial y}} = \frac{p}{1}$$
$$\frac{y + 1}{x + 4} = p \Rightarrow y = px + 4p - 1.$$

Substituting into budget constraint $px + y = 7$ we have

$$2px + 4p = 8 \Rightarrow x(p) = \frac{4 - 2p}{p} = \frac{4}{p} - 2 \Rightarrow y = 3 + 2p.$$

So buys $\frac{4}{p} - 2$ bags of crisps, $3 + 2p$ bars of chocolate.

$$\epsilon_x = p \frac{p}{4 - 2p} \left(-\frac{4}{p^2} \right) = \frac{2}{p - 2}$$
$$\Rightarrow \epsilon_x + 1 = \frac{p}{p - 2} < 0 \quad \text{if } p < 2.$$

11.

$$C(q) = q^3 - 2q^2 + 4q + 36 \Rightarrow AVC(q) = q^2 - 2q + 4.$$

$$AVC'(q) = 2q - 2 = 0 \quad \text{when} \quad q = 1 \Rightarrow \min(AVC) = 3.$$

So cease production when $p = \min(AVC) = 3$. For $p \geq 3$,

$$\begin{aligned} p = C'(q) &= 3q^2 - 4q + 4 \Rightarrow 3q^2 - 4q + 4 - p = 0 \\ \Rightarrow q &= \frac{4 \pm \sqrt{16 - 12(4 - p)}}{6} = \frac{2 \pm \sqrt{3p - 8}}{3}. \end{aligned}$$

Take +ve sign for maximum profit. So

$$S(p) = \begin{cases} \frac{2 + \sqrt{3p - 8}}{3} & \text{if } p \geq 3 \\ 0 & \text{if } p < 3. \end{cases}$$

Equilibrium is when $NS(p) = D(p)$ (N firms) so

$$\frac{2 + \sqrt{3p - 8}}{3} = 10 - p.$$

$p = 8$ is a solution by inspection, and since $D(p)$ is decreasing and $S(p)$ is increasing, it is unique.

$$\begin{aligned} p = 8 \Rightarrow q &= \frac{1}{N}D(p) = 2 \Rightarrow P(q) = pq - C(q) \\ &= 16 - (8 - 8 + 8 + 36) = -28. \end{aligned}$$

So each firm makes a loss of 28 units.

Production not viable in the long-run for $p < \min(ATC)$.

$$\begin{aligned} ATC &= q^2 - 2q + 4 + \frac{36}{q} \Rightarrow ATC'(q) = 2q - 2 - \frac{36}{q^2} \\ &= 0 \quad \text{when} \quad q = 3, \end{aligned}$$

by inspection. It is a minimum, since

$$ATC''(q) = 2 + \frac{72}{q^3} > 0.$$

$$\min(ATC) = 9 - 6 + 4 + 12 = 19.$$

So minimum price in the long-run is 19 units.

12.

$$C_1(q_1) = 6 + 7q_1 + \frac{1}{2}q_1^2,$$

$$C_2(q_2) = 4 + 8q_2 + q_2^2,$$

Profits:

$$\begin{aligned} P_1(q_1, q_2) &= pq_1 - (6 + 7q_1 + \frac{1}{2}q_1^2) = [14 - (q_1 + q_2)]q_1 - (6 + 7q_1 + \frac{1}{2}q_1^2) \\ &= -\frac{3}{2}q_1^2 - q_1q_2 + 7q_1 - 6, \end{aligned}$$

$$\begin{aligned} P_2(q_1, q_2) &= pq_2 - (4 + 8q_2 + q_2^2) = [14 - (q_1 + q_2)]q_2 - (4 + 8q_2 + q_2^2) \\ &= -q_1q_2 - 2q_2^2 + 6q_2 - 4. \end{aligned}$$

Cournot duopoly \Rightarrow maximise P_1, P_2 wrto q_1, q_2 respectively. So

$$\frac{\partial P_1}{\partial q_1} = -3q_1 - q_2 + 7 = 0,$$

$$\frac{\partial P_2}{\partial q_2} = -q_1 - 4q_2 + 6 = 0,$$

Then $q_1 = 2, q_2 = 1$. So $p = 14 - 2 - 1 = 11$ and $P_1(2, 1) = 0, P_2(2, 1) = -2$.

If co-operate, maximise

$$\begin{aligned} P(q_1, q_2) &= P_1(q_1, q_2) + P_2(q_1, q_2) \\ &= -\frac{3}{2}q_1^2 - 2q_1q_2 + 7q_1 - 2q_2^2 + 6q_2 - 10 \end{aligned}$$

$$\frac{\partial P}{\partial q_1} = -3q_1 - 2q_2 + 7 = 0,$$

$$\frac{\partial P}{\partial q_2} = -2q_1 - 4q_2 + 6 = 0,$$

giving $q_1 = 2, q_2 = \frac{1}{2}$. Then $P_1(2, \frac{1}{2}) = 1, P_2(2, \frac{1}{2}) = -\frac{5}{2}$.

13.

$$\frac{dn}{dt} = 2n^2 - 7n + 3 = (2n - 1)(n - 3) = f(n).$$

The equilibrium densities are $n = \frac{1}{2}$ and $n = 3$. The graph of $f(n)$ looks like this:

so $n = \frac{1}{2}$ is stable, $n = 3$ unstable.

Writing

$$\begin{aligned}\frac{1}{2n^2 - 7n + 3} &= \frac{A}{n - 3} + \frac{B}{2n - 1} \\ \Rightarrow 1 &= (2n - 1)A + (n - 3)B. \\ n = \frac{1}{2} &\Rightarrow B = -\frac{2}{5}, \quad n = 3 \Rightarrow A = \frac{1}{5} \\ \Rightarrow \frac{1}{5} \int_0^n \left[\frac{1}{n - 3} - \frac{2}{2n - 1} \right] &= t \\ \Rightarrow \frac{1}{5} \ln \frac{n - 3}{3(2n - 1)} &= t \\ \Rightarrow n - 3 &= 3(2n - 1)e^{5t} \Rightarrow n = 3 \frac{1 - e^{-5t}}{6 - e^{-5t}}.\end{aligned}$$

As $t \rightarrow \infty$, $n \rightarrow \frac{1}{2}$.

14.

$$\frac{dx}{dt} = x(5 - x) - xy, \quad \frac{dy}{dt} = y(7 - 2y) - xy,$$

Terms (1), (3) are logistic growth functions, implying each population could survive on its own in a limited resource environment.

Terms (2), (4) both with negative signs imply competition between the two species.

$$\begin{aligned} \text{Either } 5 - x - y = 7 - x - 2y = 0 &\Rightarrow x = 3, \quad y = 2 \\ \text{or } x = 7 - x - 2y = 0 &\Rightarrow y = \frac{7}{2}, \\ \text{or } y = 5 - x - y = 0 &\Rightarrow x = 5, \\ \text{or } x = y = 0. \end{aligned}$$

So the equilibria are $(0, 0)$, $(0, \frac{7}{2})$, $(5, 0)$, $(3, 2)$.

Community matrix

$$A = \begin{pmatrix} (5 - x - y) - x & -x \\ -y & (7 - x - 2y) - 2y \end{pmatrix}.$$

For $(0, 0)$, $A = \begin{pmatrix} 5 & 0 \\ 0 & 7 \end{pmatrix}$. E-values 5, 7 both positive \Rightarrow improper node, unstable.

For $(0, \frac{7}{2})$, $A = \begin{pmatrix} \frac{3}{2} & 0 \\ -\frac{7}{2} & -7 \end{pmatrix}$. E-values $\frac{3}{2}$, -7 opposite signs \Rightarrow saddle point.

For $(5, 0)$, $A = \begin{pmatrix} -5 & -5 \\ 0 & 2 \end{pmatrix}$. E-values -5 , 2 opposite signs \Rightarrow saddle point.

For $(3, 2)$, $A = \begin{pmatrix} -3 & -3 \\ -2 & -4 \end{pmatrix}$.

Linearised equations

$$\begin{aligned} \frac{d\epsilon_x}{dt} &= -3\epsilon_x - 3\epsilon_y \\ \frac{d\epsilon_y}{dt} &= -2\epsilon_x - 4\epsilon_y. \\ -3\epsilon_x - 3\epsilon_y &= -\frac{3}{5}\delta [3e^{-t} + 2e^{-6t}] - \frac{3}{5}\delta [-2e^{-t} + 2e^{-6t}] \\ &= -\frac{3}{5}\delta e^{-t} - \frac{12}{5}\delta e^{-6t} = \frac{d\epsilon_x}{dt} \\ -2\epsilon_x - 4\epsilon_y &= -\frac{2}{5}\delta [3e^{-t} + 2e^{-6t}] - \frac{4}{5}\delta [-2e^{-t} + 2e^{-6t}] \\ &= \frac{2}{5}\delta e^{-t} - \frac{12}{5}\delta e^{-6t} = \frac{d\epsilon_y}{dt} \end{aligned}$$

Also $\epsilon_x(0) = 1$, $\epsilon_y(0) = 0$.