

PAPER CODE NO. <b>MATH322</b>
----------------------------------



THE UNIVERSITY  
*of* LIVERPOOL

SUMMER 2000 EXAMINATIONS

Degree of Bachelor of Science : Year 3  
Degree of Master of Mathematics : Year 3  
Degree of Master of Physics : Year 3

**CHAOS THEORY**

TIME ALLOWED : Two Hours and a Half

---

INSTRUCTIONS TO CANDIDATES

Full marks can be obtained for FIVE complete answers.  
Only the best FIVE answers will be taken into account.



THE UNIVERSITY  
*of* LIVERPOOL

1. Consider the dynamical system given by iterations of a function:

$$x_{n+1} = f(x_n).$$

where the map  $f$  is  $f(x) = \mu x - x^3$ ,  $x \in \mathbb{R}$ .

- (i) Find the fixed points. Show that there is only one fixed point for  $\mu \leq 1$  and there are three for  $\mu > 1$ .

**2 marks**

- (ii) Using your results from part (i), determine stability of these fixed points for all values of the parameter  $\mu$ .

**18 marks**

2. (a) Consider the dynamical system given by iterations of a function:

$$x_{n+1} = Q(x_n).$$

where the quadratic map  $Q$  is given by

$$Q(x) = x^2 - 0.9.$$

Find the 2-periodic points and determine their stability.

**15 marks**

- (b) Consider the dynamical system given by iterations of a function  $x_{n+1} = x_n^{1/3}$ . Find the Lyapunov exponent of the map and characterise whether the behaviour of the system is chaotic or regular.

**5 marks**

3. Consider a nonlinear dynamical system defined by iterates of the functional relationships

$$x_{n+1} = f(x_n, y_n), \quad y_{n+1} = g(x_n, y_n)$$

with  $f(x, y) = 1.3x - 0.3x^2 - 0.15xy$ ,  $g(x, y) = 1.3y - 0.3y^2 - 0.15xy$ .

- (i) Find all equilibrium vectors of the system.

- (ii) Study the stability of the equilibrium vectors.

**20 marks**



THE UNIVERSITY  
*of* LIVERPOOL

4. (a) Consider an interval divided into 5 smaller equal subintervals. Consider the operation  $Q$  of removing 3 of these smaller intervals leaving 2 at the ends of the interval. Apply  $Q$  initially to the unit length interval to leave two smaller intervals. Then apply  $Q$  again to each of these smaller subintervals. The process is repeated infinitely.

Discuss whether the resulting set is self-similar under the discrete group of coordinate dilations and find its capacity dimension

**9 marks**

- (b) A dynamical system defined on  $[0, 1]$  is given by

$$\begin{aligned}x_{n+1} &= f(x_n) \quad \text{where} \\f(x) &= 4x \quad \text{for } 0 \leq x \leq \frac{1}{4} \\f(x) &= 0 \quad \text{for } \frac{1}{4} < x < \frac{3}{4} \\f(x) &= 4x - 3 \quad \text{for } \frac{3}{4} \leq x \leq 1.\end{aligned}$$

Sketch  $f(x)$ .

**1 mark**

Show that the fixed points of this system are unstable.

**2 mark**

Consider the set  $S$  of initial points  $x_0$  for which  $x_n$  is not eventually 0. Obtain a description of  $S$  and use it to find the capacity dimension of  $S$ .

**8 marks**



THE UNIVERSITY  
*of* LIVERPOOL

PAPER CODE MATH322 Page 5 of 7 END.0in

5. Consider the Weierstrass-Mandelbrot function for  $x \geq 0$

$$F(x; p) = \sum_{n=-\infty}^{\infty} p^{(D-2)n} (1 - \cos p^n x), \quad p > 1, \quad 1 < D < 2.$$

(i) Show that  $C(x; p)$  is a parametric-homogeneous function of degree  $d = (2 - D)$  and parameter  $p$ .

**3 marks**

(ii) Construct another fractal parametric-homogeneous functions  $b_2$  of degree  $d = 2$  and the same parameter  $p$ . Describe the scaling properties of the function and its trend.

**7 marks**

(iii) Assuming that the Hausdorff dimension of the graph  $C(x; p)$  is equal to its capacity dimension  $D$ , calculate the Hausdorff dimension of the  $b_2$  graph.

**5 marks**

(iv) Show that the constructed function  $b_2$  is differentiable at  $x = 0$  and find the derivative.

**5 marks**



THE UNIVERSITY  
*of* LIVERPOOL

PAPER CODE MATH322 Page 6 of 7 END.0in

6. (i) Discuss briefly some possible bifurcations that can occur as a parameter is varied in a dynamical system described by two autonomous coupled differential equations.

**2 marks**

Consider the equations describing the dynamical behaviour of a system

$$\frac{dx}{dt} = by(1 - y) - y^2x,$$

$$\frac{dy}{dt} = (1 - y)(1 - by) + y^2x$$

where  $b$  is a real positive parameter.

Find the fixed point of the system and discuss its stability for  $b > 0$ , and  $b \neq 2$  and  $b \neq 4$ .

**11 marks**

- (ii) For the particular case when  $b = 3$ , consider trajectories which pass through the four points  $(0, 1/3)$ ,  $(1, 1)$ ,  $(0, 2)$ , and  $(-1, 1)$  and sketch the directions of the tangents to these trajectories.

**5 marks**

Give plausibility arguments that the system has a stable limit cycle when  $b = 3$ .

**2 marks**



THE UNIVERSITY  
*of* LIVERPOOL

7. Consider the equations

$$\frac{dx}{dt} = -z$$

$$\frac{dy}{dt} = -x - y + 2xz$$

$$\frac{dz}{dt} = 2y - y(y - x)$$

Show that the trajectories cannot cross in phase space.

**2 marks**

Show that this system of equations describes a dissipative system.

**2 marks**

Show that the origin is a fixed point and find the other fixed point of these equations.

**4 marks**

Consider first the fixed point at the origin: linearise the equations about this fixed point and show that one eigenvalue is 1 and find the others. Discuss the stability of this fixed point.

**4 marks**

For other fixed point: linearise the equations about it and find the characteristic equations for the eigenvalues. Show that one of these eigenvalues is -2 and find the others.

**4 marks**

Without evaluating the eigenvectors, describe briefly the motion of trajectories which are near each of these two fixed points.

**4 marks**