

This paper is not to be removed from the Examination Halls

UNIVERSITY OF LONDON

279 0095 ZB

BSc degrees and Diplomas for Graduates in Economics, Management, Finance and the Social Sciences, the Diploma in Economics and Access Route for Students in the External Programme

Further Mathematics for Economists

Friday, 9 June 2006 : 10.00am to 1.00pm

Candidates should answer **EIGHT** of the following **TEN** questions: **SIX** from Section A (60 marks in total) and **TWO** from Section B (20 marks each). **Candidates are strongly advised to divide their time accordingly.**

Throughout, **R** denotes the set of real numbers.

Graph paper is provided. If used, it must be fastened securely inside the answer book.

A hand held calculator may be used when answering questions on this paper but it must not be pre-programmed or able to display graphics, text or algebraic equations. The make and type of machine must be stated clearly on the front cover of the answer book.

PLEASE TURN OVER

SECTION A

Answer all six questions from this section (60 marks in total).

- 1.(a) Show that the set $\{\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3, \mathbf{x}_4\}$ of vectors in \mathbb{R}^4 is linearly independent, where

$$\mathbf{x}_1 = \begin{pmatrix} 1 \\ 2 \\ 1 \\ 2 \end{pmatrix}, \quad \mathbf{x}_2 = \begin{pmatrix} 2 \\ -1 \\ 5 \\ 3 \end{pmatrix}, \quad \mathbf{x}_3 = \begin{pmatrix} 3 \\ 8 \\ -7 \\ 0 \end{pmatrix}, \quad \mathbf{x}_4 = \begin{pmatrix} 1 \\ 2 \\ -1 \\ 0 \end{pmatrix}.$$

- (b) Suppose that V is a vector space. What does it mean to say that the subset W of V is a 'subspace' of V ?

Suppose that V is the vector space of all 2×2 matrices with real numbers as entries, with the usual addition and scalar multiplication operations. (You may assume that this is indeed a vector space.) Show that the set of all diagonal 2×2 matrices is a subspace of V . Show, however, that the sets

$$W = \left\{ \begin{pmatrix} 1 & a \\ b & 1 \end{pmatrix} : a, b \in \mathbb{R} \right\} \quad \text{and} \quad U = \left\{ \begin{pmatrix} 0 & a^2 \\ b^2 & 0 \end{pmatrix} : a, b \in \mathbb{R} \right\}$$

are *not* subspaces of V .

2. What does it mean to say that a basis of a vector space is an 'orthonormal' basis?

Find an orthonormal basis for the subspace W of \mathbb{R}^3 given by

$$W = \left\{ \begin{pmatrix} x \\ y \\ z \end{pmatrix} : x + 2y - 3z = 0 \right\}.$$

Extend this to an orthonormal basis of \mathbb{R}^3 .

3. What, precisely, does it mean to say that a sequence (x_n) of real numbers converges to the real number L as n tends to infinity? Prove, using this precise definition, that if

$$x_n = \frac{4n^2 + 2n - 3}{2n^2 - 2n + 5},$$

then $x_n \rightarrow 2$ as $n \rightarrow \infty$.

PLEASE TURN OVER

4. Suppose A and B are non-empty bounded sets of positive real numbers, and that $C = \{a + 2b : a \in A, b \in B\}$. Prove that C is bounded above and that

$$\sup C = \sup A + 2 \sup B.$$

(Here, for a bounded non-empty set S of real numbers, $\sup S$ denotes the supremum of S .)

5. Find the value and optimal mixed strategies of the matrix game with pay-off matrix

$$\begin{pmatrix} 4 & 3 \\ 2 & 5 \end{pmatrix},$$

explaining your method.

6. By solving its dual, find the solution of the following linear programming problem:

$$\text{Minimise } 40x + 70y + 90z$$

$$\text{Subject to : } x + 2y + z \geq 4$$

$$x + y + 3z \geq 6$$

$$x, y, z \geq 0.$$

SECTION B

Answer **two** questions from this section (20 marks each).

7.(a) The function g is given by

$$g(x, y, z) = 5x^2 + y^2 + z^2 + 2xy + Axz,$$

where A is a positive constant. Show that g is convex if $A < 4$ but that it is not convex if $A > 4$.

(b) Suppose that

$$A = \begin{pmatrix} 2 & 1 & 1 \\ 1 & 1 & 2 \\ 1 & 2 & 1 \end{pmatrix}, \quad \mathbf{v} = \begin{pmatrix} 1 \\ 1 \\ 1 \end{pmatrix}.$$

Show that \mathbf{v} is an eigenvector of A , and find the corresponding eigenvalue. Find the remaining eigenvalues of A and, for each of these eigenvalues, find a corresponding eigenvector.

Find an orthogonal matrix P and a diagonal matrix D such that $P^T A P = D$.

Let f denote the quadratic form

$$f(x, y, z) = 2x^2 + y^2 + z^2 + 2xy + 2xz + 4yz.$$

Show that it is possible to write f in the form

$$f(x, y, z) = \lambda_1 X^2 + \lambda_2 Y^2 + \lambda_3 Z^2,$$

for some numbers $\lambda_1, \lambda_2, \lambda_3$, where x, y, z are linear combinations of X, Y, Z . (You should state explicitly how x, y, z may be written in terms of X, Y, Z .)

Hence, or otherwise, find a vector \mathbf{x} for which $f(x, y, z) = -9$.

PLEASE TURN OVER

- 8.(a) What is meant by a *linear* mapping T from \mathbb{R}^n to \mathbb{R}^m ? Show that, if A is any $m \times n$ matrix, then the function $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $T_A(\mathbf{x}) = A\mathbf{x}$ is linear.

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^4$ be given by

$$T \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} y \\ x \\ x + y \\ x - 2y \end{pmatrix}.$$

Write down a matrix A such that $T = T_A$. Determine the rank and nullity of T .

- (b) Find the limit of the sequence (x_n) where

$$x_n = \sum_{i=1}^n \frac{i}{n^2} = \frac{1}{n^2} + \frac{2}{n^2} + \cdots + \frac{2n}{n^2}.$$

- (c) State the Intermediate Value Theorem.

Suppose the continuous function $f : \mathbb{R} \rightarrow \mathbb{R}$ is not bounded above and not bounded below. Prove that $\{f(x) : x \in \mathbb{R}\} = \mathbb{R}$.

- 9.(a) A sequence (x_n) is defined as follows: $x_1 = 1$ and, for $n \geq 1$,

$$x_{n+1} = \frac{x_n + 1}{4}.$$

Prove that, for all $n \in \mathbb{N}$, $x_n > 1/3$. Prove also that for all $n \in \mathbb{N}$, $x_{n+1} \leq x_n$. Why does it follow that the sequence converges? Determine the limit of x_n as n tends to infinity.

- (b) What is meant by an ‘open’ subset of \mathbb{R}^n ? What is meant by a ‘closed’ subset of \mathbb{R}^n ? State a result that characterises closed sets in terms of open sets.

Prove that the subset

$$U = \left\{ \begin{pmatrix} x \\ y \end{pmatrix} : x + y > 0 \right\}$$

is an open subset of \mathbb{R}^2 .

10.(a) Consider the functions

$$f(x, y, z) = x + \frac{4}{y} - \frac{2y}{x} + 6z - z^3 \quad (x \neq 0, y \neq 0)$$

$$g(x, y, z) = x^2 - yz + y^2 + z^2 - 3x - 2y.$$

Show that the point $P = (2, 2, 1)$ is on both of the surfaces

$$f(x, y, z) = 7, \quad g(x, y, z) = -3.$$

Find the equations of the tangent planes to each of these surfaces at the point $(2, 2, 1)$.

Show that these two planes are orthogonal.

For each function, find the critical (or stationary) points and determine the nature of each critical point.

(b) Two consumers engaged in a barter arrangement, R and M , have utility functions $u_R(x, y)$ and $u_M(x, y)$, respectively, for bundles (x, y) of x bushels of wheat and y bushels of oats, where

$$u_R(x, y) = x^{1/3}y, \quad u_M(x, y) = x^2y^{1/2}.$$

Given that they have 20 bushels of wheat in total and 10 bushels of oats in total, determine the equation of the contract curve.

END OF PAPER

