

1999 Question 1

(a) 2 marks

$$(1 + i)^4 = (1 + 2i - 1)^2 = (2i)^2 = -4$$

(b) 3 marks

$$\cos(\pi - i \log_e 2) = \cos \pi \cos(i \log_e 2) + \sin \pi \sin(i \log_e 2)$$

(Unit A2, Section 4, Para. 5)

$$= -\cos(i \log_e 2)$$

$$= -\frac{\exp(-\log_e 2) + \exp(\log_e 2)}{2} \quad (\text{Unit A2, Section 4, Para. 4})$$

$$= -\frac{\frac{1}{2} + 2}{2} = -\frac{5}{4}$$

(c) 3 marks

$$(-e)^{i\pi} = \exp(i\pi \operatorname{Log}(-e)) \quad (\text{Unit A2, Section 5, Para. 3})$$

$$= \exp(i\pi (\log_e |-e| + i \operatorname{Arg}(-e))) \quad (\text{Unit A2, Section 5, Para. 1})$$


$$= \exp(i\pi (1 + i\pi))$$

$$= \exp(i\pi) \exp(-\pi^2)$$

$$= -\exp(-\pi^2)$$


1999 Question 2

(a) 2 marks



(b) 6 marks

$C = \text{ext } A$ (Unit A3, Section 5, Para. 9)



(b)(i) (Unit A3, Section 4, Paras. 6 and 7)

A is not a region since it is not open.
 B is a region.
 C is not a region as it is not connected.

(b)(ii) (Unit A3, Section 5, Para. 5)

A is compact.
 B is not compact as it is not closed or bounded.
 C is not compact as it is not closed or bounded.

1999 Question 3

(a) 4 marks (Unit B1, Ex. 2.1(ii) – Opposite direction along contour)

The standard parametrization (Unit A2, Section 2, Para. 3) for Γ is

$$\gamma(t) = (1-t)i + t, \quad t \in [0, 1]$$

and $\gamma'(t) = 1 - i$.

As γ is differentiable on $[0, 1]$, γ' is continuous on $[0, 1]$, and γ' is non-zero on $[0, 1]$ then γ is a smooth path (Unit A4, Section 4, Para. 3).

As γ is a smooth parametrization (Unit B1, Section 2, Para. 1) then

$$\begin{aligned} \int_{\Gamma} \operatorname{Im} z \, dz &= \int_0^1 \{ \operatorname{Im} \gamma(t) \} \gamma'(t) \, dt \\ &= \int_0^1 (1-t)(1-i) \, dt \\ &= (1-i) \left[-\frac{(1-t)^2}{2} \right]_0^1 \\ &= \frac{1-i}{2} \end{aligned}$$

(b) 4 marks

The length of C is $L = 2\pi * 2 = 4\pi$.

Using the Triangle Inequality (Unit A1, Section 5, Para. 3b) then, for $z \in C$, we have

$$|\bar{z}^2 - 1| \leq |\bar{z}^2| + 1 = |z|^2 + 1 = 4 + 1 = 5$$

Using the Backwards form of the Triangle Inequality (Unit A1, Section 5, Para. 2b) then, for $z \in C$, we have

$$|z^2 - 1| \geq \left| |z^2| - 1 \right| = |4 - 1| = 3$$

Putting $f(z) = \frac{\bar{z}^2 - 1}{z^2 - 1}$ we have $|f(z)| \leq 5/3 = M$ for $z \in C$.

By the Quotient Rule (Unit A3, Section 2, Para. 5) $f(z)$ is continuous on $\mathbb{C} - \{-1, 1\}$ and hence on the circle C .

Therefore by the Estimation Theorem (Unit B1, Section 4, Para. 3)

$$\left| \int_{\Gamma} f(z) \, dz \right| \leq ML = \frac{5}{3} * 4\pi = \frac{20}{3} \pi$$

1999 Question 4

(a) 6 marks

(a)(i) \mathbb{C} is a simply-connected region (Unit B2, Section 1, Para. 3), C is a simple-closed contour (Unit B2, Section 1, Para. 1) in \mathbb{C} , and $f(z) = \exp(i\pi z)$ is analytic on \mathbb{C} .

As -1 lies inside the circle C then by Cauchy's Integral formula (Unit B2, Section 2, Para. 1)

$$\int_C \frac{e^{i\pi z}}{z+1} dz = 2\pi i f(-1) = 2\pi i * e^{-i\pi} = -2\pi i$$

(a)(ii) Let $R = \{z \in \mathbb{C} : |z - i| < 5^{1/2}\}$. R is a simply-connected region (Unit B2, Section 1, Para. 3) and C is a simple-closed contour in R . As $\frac{e^{i\pi z}}{z+2}$ is analytic on R then by Cauchy's Theorem (Unit B2, Section 1, Para. 4)

$$\int_C \frac{e^{i\pi z}}{z+2} dz = 0$$

(b) 2 marks

Since $1 + z^2$ and \cos are entire functions then by the Composition rule (Unit A4, Section 3, Para. 1) so is $f(z) = \cos(1 + z^2)$.

By Liouville's theorem (Unit B2, Section 2, Para. 2) if f is a bounded entire function then f is constant.

Since $f(i) = \cos 0 = 1 \neq \cos 1 = f(0)$ then f is not constant. Therefore f is not bounded on \mathbb{C} so there is a complex number z such that $|\cos(1 + z^2)| > 100$.

1999 Question 5

Same as 2004 Qu 5 apart from 4 marks being awarded for each part.

1999 Question 6

Same as 2004 Qu 6.


1999 Question 7

Equivalent to 2004 Qu 7 except in part (b) also need to calculate the stream line through i .

A streamline through i is given by $\frac{1}{2}(-x^2 + y^2) = \Psi(0,1) = \frac{1}{2}$.

Therefore the streamline has the equation $y^2 = x^2 + 1$.

At i the velocity function $q(i) = i(-i) = 1$ (positive x direction)



1999 Question 8

(a) 3 marks

Since $f(i) = i^2 - i^2 + i = i$ then i satisfies the fixed point equation (Unit D3, Section 1, Para. 3). Therefore i is a fixed point of f .

$$f'(z) = 2z - i.$$

As $|f'(i)| = |i| = 1$ then i is an indifferent fixed point of f . (Unit D3, Section 1, Para. 5)

(b) 5 marks

(b)(i) [Unit D3, Problem 4.3(b)]

Let $c = 1 + i$.

$$P_c(0) = 1 + i.$$

$$P_c^2(0) = (1 + i)^2 + (1 + i) = 2i + (1 + i) = 1 + 3i.$$

As $|P_c^2(0)| > 2$ then c does not lie in the Mandelbrot set (Unit D3, Section 4, Para. 5).

(b)(ii) Let $c = -\frac{9}{10} - \frac{\sqrt{3}}{10}i$.

Since $|c + 1| = \left| \frac{1}{10} - \frac{\sqrt{3}}{10}i \right| = \frac{1}{5} < \frac{1}{4}$ then P_c has an attracting 2-cycle (Unit D3, Section 4, Para. 9).

Therefore c belongs to the Mandelbrot set (Unit D3, Section 4, Para. 8).

1999 Question 9

(a) 8 marks

(a)(i)

$$f(z) = z + |z|^2 = (x + iy) + (x^2 + y^2) = u(x, y) + iv(x, y),$$

where $u(x, y) = x + x^2 + y^2$, and $v(x, y) = y$.

$$\frac{\partial u}{\partial x}(x, y) = 1 + 2x, \quad \frac{\partial u}{\partial y}(x, y) = 2y, \quad \frac{\partial v}{\partial x}(x, y) = 0, \quad \frac{\partial v}{\partial y}(x, y) = 1$$

(a)(ii)

If f is differentiable then the Cauchy-Riemann equations hold (Unit A4, Section 2, Para. 1). If they hold at (a, b)

$$\frac{\partial u}{\partial x}(a, b) = 1 + 2a = 1 = \frac{\partial v}{\partial y}(a, b), \text{ and}$$

$$\frac{\partial v}{\partial x}(a, b) = 0 = -2b = -\frac{\partial u}{\partial y}(a, b)$$

Therefore the Cauchy-Riemann equations only hold at $(0, 0)$.

As f is defined on the region \mathbb{C} , and the partial derivatives $\frac{\partial u}{\partial x}, \frac{\partial u}{\partial y}, \frac{\partial v}{\partial x}, \frac{\partial v}{\partial y}$

1. exist on \mathbb{C}
2. are continuous at $(0, 0)$.
3. satisfy the Cauchy-Riemann equations at $(0, 0)$

then, by the Cauchy-Riemann Converse Theorem (Unit A4, Section 2, Para. 3), f is differentiable at 0 .

As the Cauchy-Riemann only hold at $(0, 0)$ then f is not differentiable on any region surrounding 0 . Therefore f is not analytic at 0 . (Unit A4, Section 1, Para. 3)


(a)(iii)

$$f'(0, 0) = \frac{\partial u}{\partial x}(0, 0) + i \frac{\partial v}{\partial x}(0, 0) = 1 \quad (\text{Unit A4, Section 2, Para. 3}).$$

(b) 10 marks


(i) The domain of g is \mathbb{C} (Unit A4, Section 1, Para. 7) and its derivative $g'(z)=3z^2$ also has domain \mathbb{C} (Unit A4, Section 3, Para. 4). Therefore g is analytic at i and since $g'(i) = -3 \neq 0$ then g is conformal at i (Unit A4, Section 4, Para. 6).

(ii) $\pi/2$ is in the domain of γ_1 so $\gamma_1(\pi/2) = e^{i\pi/2} = i$.
 1 belongs to the domain of γ_2 so $\gamma_2(1) = i$. Therefore Γ_1 and Γ_2 meet at the point i .



(iii) As g is analytic on \mathbb{C} and $g'(i) \neq 0$ then a small disc centred at i is mapped approximately (Unit A4, Section 1, Para. 11) to a small disc centred at $g(i) = -i$. The disc is rotated by $\text{Arg}(g'(i)) = \text{Arg}(-3) = \pi$, and scaled by a factor $|g'(i)| = 3$.

The horizontal line in the diagram below is $g(\Gamma_1)$. (Unit A4, Section 4, Para. 4)



1999 Question 10

(a) 7 marks

$$\sin z = z \left(1 - \frac{z^2}{3!} + \frac{z^4}{5!} - \dots \right), \text{ for } z \in \mathbf{C}. \quad (\text{Unit B3, Section 3, Para. 5})$$

$$\text{Since } 0 < \frac{\sin z}{z} < 1 \text{ when } 0 < |z| < \pi \text{ then } \left| -\frac{z^2}{3!} + \frac{z^4}{5!} - \dots \right| < 1.$$

$$\begin{aligned} \frac{z}{\sin z} &= \left(1 - \frac{z^2}{3!} + \frac{z^4}{5!} - \dots \right)^{-1} \\ &= 1 + \left(\frac{z^2}{3!} - \frac{z^4}{5!} + \dots \right) + \left(\frac{z^2}{3!} - \frac{z^4}{5!} + \dots \right)^2 + \dots \\ &= 1 + \frac{z^2}{6} + z^4 \left(-\frac{1}{120} + \frac{1}{36} \right) + \dots \end{aligned}$$

Therefore the Laurent series about 0 for f is $1 + \frac{z^2}{6} + \frac{7}{360}z^4 + \dots$ for $0 < |z| < \pi$

$\frac{1}{z^2 \sin z} = \frac{f(z)}{z^3}$ is analytic on the punctured disc $\mathbf{C} - \{0\}$. It has the Laurent series about 0

$$\frac{1}{z^3} + \frac{1}{6z} + \frac{7}{360}z + \dots = \sum_{n=-\infty}^{\infty} a_n z^n$$

As C is a circle with centre 0 then (Unit B4, Section 4, Para. 2)

$$\int_C \frac{f(z)}{z^3} dz = 2\pi i a_{-1} = \frac{\pi i}{3}.$$

(b) 5 marks

The domain of $A = \mathbf{C} - \{n\pi : n \in \mathbf{Z}\}$.

Suppose that g is another analytic function with domain A which agrees with f on $\{iy : y > 0\}$

The set $S = \{i(1 + \frac{1}{n}) : n = 1, 2, 3, \dots\} \subseteq A$ and has the limit point $i \in A$.

f agrees with g throughout the set $S \subseteq A$ and S has a limit point which is in A . Therefore by the Uniqueness theorem (Unit B3, Section 5, Para. 7) f agrees with g throughout A . Hence f is the only analytic function with domain A such that $f(iy) = \frac{y}{\sinh y}$ for $y > 0$.

(c) 6 marks

Since $\sin z = 0$ when $z = 0, z = \pm\pi, z = \pm 2\pi, \dots$. Then $f(z)$ has singularities of the form $k\pi, k \in \mathbb{Z}$.

Singularity at $z = 0$.

At $z = 0$ we can use the Laurent series found in part (a). Since $f(0) = 1$ then the singularity at 0 is a removable singularity.

Singularities at $z = k\pi$ where $k \in \mathbb{Z} - \{0\}$.

$$\sin(z - k\pi) = \sin z * \cos k\pi - \cos z * \sin k\pi = (-1)^k \sin z.$$

$$\text{Therefore } f(z) = \frac{z}{\sin z} = (-1)^k \frac{z}{\sin(z - k\pi)} = (-1)^k \left\{ \frac{z - k\pi}{\sin(z - k\pi)} + \frac{k\pi}{\sin(z - k\pi)} \right\}$$

$$\lim_{z \rightarrow k\pi} \frac{z - k\pi}{\sin(z - k\pi)} = 1$$

As $\lim_{z \rightarrow k\pi} (z - k\pi) \frac{k\pi}{\sin(z - k\pi)} = k\pi$ then there is a simple pole at $z = k\pi$ (Unit B4, Section 3, Para. 2).

Therefore there f has simple poles at $k\pi$ where $k \in \mathbb{Z} - \{0\}$.

1999 Question 11

(a) 6 marks

Since $f(z) = \frac{\pi \cot \pi z}{9(z - i/3)(z + i/3)}$ then f has simple poles at $z = \pm i/3$.

By the cover-up rule (Unit C1, Section 1, Para. 3)

$$\operatorname{Res}(f, i/3) = \frac{\pi \cot(i\pi/3)}{9(i/3 + i/3)} = \frac{\pi \cot(i\pi/3)}{6i}, \text{ and}$$

$$\operatorname{Res}(f, -i/3) = \frac{\pi \cot(-i\pi/3)}{9(-i/3 - i/3)} = \frac{\pi \cot(-i\pi/3)}{-6i}.$$

Since $\sin(iz) = i \sinh z$ and $\cos(iz) = \cosh z$ then $\cot(iz) = -i \coth(z)$.

Therefore $\operatorname{Res}(f, i/3) = -\frac{\pi \coth(\pi/3)}{6}$ and

$$\operatorname{Res}(f, -i/3) = \frac{\pi \coth(-\pi/3)}{6} = -\frac{\pi \coth(\pi/3)}{6}. \text{ (Unit A2, Section 4, Para. 6)}$$

$f(z) = g(z) / h(z)$ where $g(z) = \frac{\pi \cos \pi z}{9z^2 + 1}$ and $h(z) = \sin \pi z$.

g and h are analytic at 0, $h(0) = 0$, and $h'(0) = \pi \cos(0) = \pi \neq 0$.

Therefore by the g/h rule (Unit C1, Section 1, Para. 2)

$$\operatorname{Res}(f, 0) = \frac{g(0)}{h'(0)} = \frac{\pi * 1}{1} * \frac{1}{\pi} = 1.$$

[You could also use Unit C1, Section 4, Para 1 – last line]

(b) 8 marks

The method given in Unit C1, Section 4, Para. 1 will be used.

$$f(z) = \pi \cot \pi z * \phi(z) \text{ where } \phi(z) = 1/(9z^2 + 1).$$

ϕ is an even function which is analytic on \mathbb{C} except for simple poles at the non-integral points $z = \pm i/3$.

Let S_N be the square contour with vertices at $(N + \frac{1}{2})(\pm 1 \pm i)$.

On S_N we have $|z| \geq N + \frac{1}{2}$ so, using the backwards form of the Triangle Inequality (Unit A1, Section 5, Para. 2),

$$|9z^2 + 1| \geq |9z^2| - 1 \geq 9(N + \frac{1}{2})^2 - 1 \geq 9N^2.$$

On S_N we also have $\cot \pi z \leq 2$ (Unit C1, Section 4, Para. 2) so on C_N

$$|f(z)| \leq \frac{\pi(2)}{9N^2}.$$

The length of the contour S_N is $4(2N + 1)$.

As f is continuous on the contour S_N then by the Estimation Theorem (Unit B1, Section 4, Para. 3) we have

$$\left| \int_{S_N} f(z) dz \right| \leq \frac{2\pi}{9N^2} 4(2N + 1) = \frac{8\pi}{9N} \left(2 + \frac{1}{N}\right).$$

$$\text{Hence } \lim_{N \rightarrow \infty} \left| \int_{S_N} f(z) dz \right| = 0.$$

Therefore the conditions specified in Unit C1, Section 4, Para. 1 hold so

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{1}{9n^2 + 1} &= -\frac{1}{2} (\text{Res}(f, 0) + \text{Res}(f, i/3) + \text{Res}(f, -i/3)) \\ &= -\frac{1}{2} + \frac{\pi}{6} \coth \frac{\pi}{3}. \end{aligned}$$

(c) 4 marks

$$\begin{aligned} \sum_{n=-\infty}^{\infty} \frac{1}{9n^2 + 1} &= \sum_{n=-\infty}^{-1} \frac{1}{9n^2 + 1} + 1 + \sum_{n=1}^{\infty} \frac{1}{9n^2 + 1} \\ &= 1 + 2 \sum_{n=1}^{\infty} \frac{1}{9n^2 + 1} = \frac{\pi}{3} \coth \frac{\pi}{3}. \end{aligned}$$

1999 Question 12

(a) 3 marks


Using the formula for a transformation mapping points to the standard triple (Unit D1, Section 2, Para. 11) then \hat{f}_1 corresponds to

$$f_1(z) = \frac{(z-0)(\infty-(1+i))}{(z-(1+i))(\infty-0)} = \frac{z}{z-(1+i)}$$

$$\text{So } \hat{f}_1\left(\frac{1}{2}(1+i)\right) = \frac{\frac{1}{2}(1+i)}{\frac{1}{2}(1+i)-(1+i)} = -1$$

(b) 15 marks

(b)(i)



(b)(ii)

As f_1 is a Möbius transformation and R is a region in the domain of f_1 then $f_1(R)$ is a region and the boundary of R maps onto the boundary of $f_1(R)$ (Unit D1, Section 4, Para. 3).

The boundary of R is $B_1 \cup B_2$ where

$$B_1 = \{z : |z-1| = 1, \pi/2 \leq \text{Arg}(z-1) \leq \pi\} \text{ and}$$

$$B_2 = \{z : |z-i| = 1, -\pi/2 \leq \text{Arg}(z-i) \leq 0\}.$$

As \hat{f}_1 maps 0 to 0 and $1+i$ to ∞ then both B_1 and B_2 are mapped to extended lines which start at the origin. Since the angle between B_1 and B_2 at 0 is $\pi/2$ and the transformation \hat{f}_1 is conformal, then the angle between the extended lines at the origin is also $\pi/2$.

Since the angle between the boundaries of R at 0 and a line joining the origin to

$\frac{1}{2}(1+i)$ is $\pi/4$ then as \hat{f}_1 is conformal then this is also true in $\hat{f}_1(R)$. From part (a) we have

$\hat{f}_1\left(\frac{1}{2}(1+i)\right) = -1$ so the 2 boundaries must be mapped to extended lines at angles $\pi/4$ above and below the negative real-axis.

Therefore $\hat{f}_1(\mathbb{R}) = S$.

(b)(iii)

$w = g(z) = z \exp(-3\pi i / 4)$ maps S to T .

Therefore the extended conformal mapping from \mathbb{R} to T is \hat{f} , where

$$\begin{aligned} f(z) &= (g \circ f_1)(z) = \frac{z \exp(-3\pi i / 4)}{z - (1+i)} = \frac{-z \frac{1}{\sqrt{2}}(1+i)}{z - (1+i)} \\ &= \frac{-\sqrt{2}z}{(1-i)z - 2} \end{aligned}$$

(b)(iv)

$$\text{Let } f(z) = \frac{-\sqrt{2}z}{(1-i)z - 2} = \frac{az + b}{cz + d}$$

Using the formula for the inverse function (Unit D1, Section 2, Para. 6) we have

$$f^{-1}(z) = \frac{-2z}{-(1-i)z - \sqrt{2}} = \frac{2z}{(1-i)z + \sqrt{2}}$$