

FUNDAMENTALS OF CHEMISTRY 1B - CHEM1002

SECOND SEMESTER EXAMINATION

CONFIDENTIAL

NOVEMBER 2007

TIME ALLOWED: THREE HOURS

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

FAMILY NAME		SID NUMBER	
OTHER NAMES		TABLE NUMBER	

INSTRUCTIONS TO CANDIDATES

- All questions are to be attempted. There are 18 pages of examinable material.
- Complete the written section of the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new question of the short answer section begins with a •.
- Electronic calculators, including programmable calculators, may be used. Students are warned, however, that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution.
- Numerical values required for any question, standard electrode reduction potentials, a Periodic Table and some useful formulas may be found on the separate data sheet.
- Pages 14, 16, 19, 23 and 24 are for rough working only.

OFFICIAL USE ONLY

Multiple choice section

	Marks	
Pages	Max	Gained
2-10	36	

Short answer section

Page	Marks		Marker
	Max	Gained	
11	8		
12	4		
13	8		
15	10		
17	8		
18	7		
20	6		
21	7		
22	6		
Total	64		

Marks
3

- Define what is meant by an “allotrope”. Give an example of a *pair* of allotropes involving (i) carbon and (ii) oxygen.


5

- The K_{sp} of $\text{Fe}(\text{OH})_3$ is $2.0 \times 10^{-39} \text{ M}^4$. What is the solubility of $\text{Fe}(\text{OH})_3$ in g L^{-1} ?

Answer:

What effect does lowering the pH have on the solubility of $\text{Fe}(\text{OH})_3$? Explain your answer.

- | |
|--------------|
| Marks |
| 4 |



RATE LAW	RATE CONSTANT
Answer:	Answer:

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

Marks
8

- Solution A consists of a 0.50 M aqueous solution of HF at 25 °C. Calculate the pH of Solution A. The pK_a of HF is 3.17.

pH =

At 25 °C, 1.00 L of Solution B consists of 12.97 g of lithium fluoride, LiF, dissolved in water. Calculate the pH of Solution B.

pH =

Solution B (1.00 L) is poured into Solution A (1.00 L) and allowed to equilibrate at 25 °C. Calculate the pH of the final solution.

pH =

If you wanted to adjust the pH of the mixture of Solution A and Solution B to be exactly equal to 4.00, which component in the mixture would you need to increase in concentration?

Marks
2

- Briefly explain why H_2S is a stronger Brønsted acid than H_2O .

2

- Compounds of d -block elements are frequently paramagnetic. Using the box notation to represent atomic orbitals, account for this property in compounds of Ni^{2+} .

6






- Complete the following table.

Formula	Oxidation state of transition metal	Coordination number of transition metal	Number of d -electrons in the complex ion	Species formed upon dissolving in water
$\text{K}_3[\text{Mn}(\text{CN})_6]$				
$[\text{Ru}(\text{NH}_3)_5(\text{OH}_2)](\text{NO}_3)_2$				
$[\text{Cr}(\text{en})_3]\text{Cl}_3$				

en = ethylenediamine = $\text{NH}_2\text{CH}_2\text{CH}_2\text{NH}_2$



Marks
8

- Give the name of the starting material where indicated and the constitutional formula of the major organic product formed in each of the following reactions.

**Name:****Name:****Name:**

Marks
4

- Classify the starting materials for each of the following reactions as nucleophile or electrophile in the boxes provided and indicate with δ^+ and δ^- the polarisation of the H–Br and C–Br bonds in the starting materials.


3

- Draw the constitutional formula for each of the following compounds.

(Z)-4-methylhex-2-ene


trans-1,3-dichlorocyclohexane

(*R*)-butan-2-ol

Marks

6

- Consider the following reaction sequence.




Give the reagents **B** and **D** and draw the structures of the major organic products, **A**, **C**, **E** and **F**, formed in these reactions.

A	D
B	E
C	F

Marks
7

- Dopa is a non-proteinogenic amino acid used to treat Parkinson's disease. Only the enantiomer (**X**) is effective in restoring nerve function. The other enantiomer is highly toxic.



What is the molecular formula of (**X**)?

List the substituents attached to the stereogenic centre in descending order of priority according to the sequence rules.

highest priority		lowest priority	

What is the absolute stereochemistry of (**X**)? Write (*R*) or (*S*).


Name the functional groups, highlighted by the boxes **a** and **b**, present in (**X**).

a =	b =
------------	------------

Give the constitutional formula of the product obtained when (**X**) is treated with NaHCO_3 .

Marks
6

- Show clearly the reagents you would use to carry out the following chemical conversion. Exactly one intermediate compound and hence two steps are required. Give the constitutional formula of the intermediate compound.



How could you distinguish between the starting material, the intermediate compound and the final product using infrared spectroscopy?

CHEM1002 - CHEMISTRY 1B**DATA SHEET***Physical constants*Avogadro constant, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$ Faraday constant, $F = 96485 \text{ C mol}^{-1}$ Planck constant, $h = 6.626 \times 10^{-34} \text{ J s}$ Speed of light in vacuum, $c = 2.998 \times 10^8 \text{ m s}^{-1}$ Rydberg constant, $E_R = 2.18 \times 10^{-18} \text{ J}$ Boltzmann constant, $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$ Gas constant, $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$ Charge of electron, $e = 1.602 \times 10^{-19} \text{ C}$ Mass of electron, $m_e = 9.1094 \times 10^{-31} \text{ kg}$ Mass of proton, $m_p = 1.6726 \times 10^{-27} \text{ kg}$ Mass of neutron, $m_n = 1.6749 \times 10^{-27} \text{ kg}$ *Properties of matter*

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm^{-3} *Conversion factors*

1 atm = 760 mmHg = 101.3 kPa

0 °C = 273 K

1 L = 10^{-3} m^3 1 Å = 10^{-10} m 1 eV = $1.602 \times 10^{-19} \text{ J}$ 1 Ci = $3.70 \times 10^{10} \text{ Bq}$ 1 Hz = 1 s^{-1} *Decimal fractions*

Fraction	Prefix	Symbol
10^{-3}	milli	m
10^{-6}	micro	μ
10^{-9}	nano	n
10^{-12}	pico	p

Decimal multiples

Multiple	Prefix	Symbol
10^3	kilo	k
10^6	mega	M
10^9	giga	G

CHEM1002 - CHEMISTRY 1B**Standard Reduction Potentials, E°**

Reaction	E° / V
$\text{S}_2\text{O}_8^{2-} + 2\text{e}^- \rightarrow 2\text{SO}_4^{2-}$	+2.01
$\text{Co}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Au}(\text{s})$	+1.50
$\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2 + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{Br}_2 + 2\text{e}^- \rightarrow 2\text{Br}^-(\text{aq})$	+1.10
$\text{MnO}_2(\text{s}) + 4\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{Mn}^{3+} + 2\text{H}_2\text{O}$	+0.96
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Co}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Co}(\text{s})$	-0.28
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Li}^+(\text{aq}) + \text{e}^- \rightarrow \text{Li}(\text{s})$	-3.04

CHEM1002 - CHEMISTRY 1B*Useful formulas*

Quantum Chemistry $E = h\nu = hc/\lambda$ $\lambda = h/mv$ $4.5k_B T = hc/\lambda$ $E = -Z^2 E_R (1/n^2)$ $\Delta x \cdot \Delta(mv) \geq h/4\pi$ $q = 4\pi r^2 \times 5.67 \times 10^{-8} \times T^4$	Electrochemistry $\Delta G^\circ = -nFE^\circ$ <i>Moles of e^- = It/F</i> $E = E^\circ - (RT/nF) \times 2.303 \log Q$ $= E^\circ - (RT/nF) \times \ln Q$ $E^\circ = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^\circ - \frac{0.0592}{n} \log Q$ (at 25 °C)
Acids and Bases $pK_w = pH + pOH = 14.00$ $pK_w = pK_a + pK_b = 14.00$ $pH = pK_a + \log \{ [A^-] / [HA] \}$	Gas Laws $PV = nRT$ $(P + n^2 a/V^2)(V - nb) = nRT$
Colligative properties $\pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^\circ_{\text{solvent}}$ $p = kc$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$	Kinetics $t_{1/2} = \ln 2/k$ $k = Ae^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$
Radioactivity $t_{1/2} = \ln 2/\lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t) \text{ years}$	Thermodynamics & Equilibrium $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ $\Delta G = \Delta G^\circ + RT \ln Q$ $\Delta G^\circ = -RT \ln K$ $K_p = K_c (RT)^{\Delta n}$
Miscellaneous $A = -\log_{10} \frac{I}{I_0}$ $A = \epsilon cl$ $E = -A \frac{e^2}{4\pi\epsilon_0 r} N_A$	Mathematics If $ax^2 + bx + c = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $\ln x = 2.303 \log x$

PERIODIC TABLE OF THE ELEMENTS

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1 HYDROGEN H 1.008																	2 HELIUM He 4.003
3 LITHIUM Li 6.941	4 BERYLLIUM Be 9.012											5 BORON B 10.81	6 CARBON C 12.01	7 NITROGEN N 14.01	8 OXYGEN O 16.00	9 FLUORINE F 19.00	10 NEON Ne 20.18
11 SODIUM Na 22.99	12 MAGNESIUM Mg 24.31											13 ALUMINIUM Al 26.98	14 SILICON Si 28.09	15 PHOSPHORUS P 30.97	16 SULFUR S 32.07	17 CHLORINE Cl 35.45	18 ARGON Ar 39.95
19 POTASSIUM K 39.10	20 CALCIUM Ca 40.08	21 SCANDIUM Sc 44.96	22 TITANIUM Ti 47.88	23 VANADIUM V 50.94	24 CHROMIUM Cr 52.00	25 MANGANESE Mn 54.94	26 IRON Fe 55.85	27 COBALT Co 58.93	28 NICKEL Ni 58.69	29 COPPER Cu 63.55	30 ZINC Zn 65.39	31 GALLIUM Ga 69.72	32 GERMANIUM Ge 72.59	33 ARSENIC As 74.92	34 SELENIUM Se 78.96	35 BROMINE Br 79.90	36 KRYPTON Kr 83.80
37 RUBIDIUM Rb 85.47	38 STRONTIUM Sr 87.62	39 YTTRIUM Y 88.91	40 ZIRCONIUM Zr 91.22	41 NIOBIUM Nb 92.91	42 MOLYBDENUM Mo 95.94	43 TECHNETIUM Tc [98.91]	44 RUTHENIUM Ru 101.07	45 RHODIUM Rh 102.91	46 PALLADIUM Pd 106.4	47 SILVER Ag 107.87	48 CADMIUM Cd 112.40	49 INDIUM In 114.82	50 TIN Sn 118.69	51 ANTIMONY Sb 121.75	52 TELLURIUM Te 127.60	53 IODINE I 126.90	54 XENON Xe 131.30
55 CAESIUM Cs 132.91	56 BARIUM Ba 137.34	57-71	72 HAFNIUM Hf 178.49	73 TANTALUM Ta 180.95	74 TUNGSTEN W 183.85	75 RHENIUM Re 186.2	76 OSMIUM Os 190.2	77 IRIDIUM Ir 192.22	78 PLATINUM Pt 195.09	79 GOLD Au 196.97	80 MERCURY Hg 200.59	81 THALLIUM Tl 204.37	82 LEAD Pb 207.2	83 BISMUTH Bi 208.98	84 POLONIUM Po [210.0]	85 ASTATINE At [210.0]	86 RADON Rn [222.0]
87 FRANCIUM Fr [223.0]	88 RADIUM Ra [226.0]	89-103	104 RUTHERFORDIUM Rf [261]	105 DUBNIUM Db [262]	106 SEABORGIUM Sg [266]	107 BOHRIUM Bh [262]	108 HASSIUM Hs [265]	109 MEITNERIUM Mt [266]	110 DARMSTADTIUM Ds [271]	111 ROENTGENIUM Rg [272]							

LANTHANIDES

57 LANTHANUM La 138.91	58 CERIUM Ce 140.12	59 PRASEODYMIUM Pr 140.91	60 NEODYMIUM Nd 144.24	61 PROMETHIUM Pm [144.9]	62 SAMARIUM Sm 150.4	63 EUROPIUM Eu 151.96	64 GADOLINIUM Gd 157.25	65 TERBIUM Tb 158.93	66 DYSPROSIUM Dy 162.50	67 HOLMIUM Ho 164.93	68 ERBIUM Er 167.26	69 THULIUM Tm 168.93	70 YTTERBIUM Yb 173.04	71 LUTETIUM Lu 174.97
89 ACTINIUM Ac [227.0]	90 THORIUM Th 232.04	91 PROTACTINIUM Pa [231.0]	92 URANIUM U 238.03	93 NEPTUNIUM Np [237.0]	94 PLUTONIUM Pu [239.1]	95 AMERICIUM Am [243.1]	96 CURIUM Cm [247.1]	97 BERKELLIUM Bk [247.1]	98 CALIFORNIUM Cf [252.1]	99 EINSTEINIUM Es [252.1]	100 FERMIUM Fm [257.1]	101 MENDELEVIUM Md [256.1]	102 NOBELIUM No [259.1]	103 LAWRENCIUM Lr [260.1]

ACTINIDES