

FUNDAMENTALS OF CHEMISTRY 1B - CHEM1002

SECOND SEMESTER EXAMINATION

CONFIDENTIAL

NOVEMBER 2008

TIME ALLOWED: THREE HOURS

GIVE THE FOLLOWING INFORMATION IN BLOCK LETTERS

FAMILY NAME		SID NUMBER	
OTHER NAMES		TABLE NUMBER	

INSTRUCTIONS TO CANDIDATES

- All questions are to be attempted. There are 19 pages of examinable material.
- Complete the written section of the examination paper in **INK**.
- Read each question carefully. Report the appropriate answer and show all relevant working in the space provided.
- The total score for this paper is 100. The possible score per page is shown in the adjacent tables.
- Each new question of the short answer section begins with a •.
- Electronic calculators, including programmable calculators, may be used. Students are warned, however, that credit may not be given, even for a correct answer, where there is insufficient evidence of the working required to obtain the solution.
- Numerical values required for any question, standard electrode reduction potentials, a Periodic Table and some useful formulas may be found on the separate data sheet.
- Pages 14, 16, 20 and 24 are for rough working only.

OFFICIAL USE ONLY

Multiple choice section

	Marks	
Pages	Max	Gained
2-9	32	

Short answer section

Page	Marks		Marker
	Max	Gained	
10	8		
11	6		
12	4		
13	6		
15	8		
17	8		
18	6		
19	6		
21	5		
22	7		
23	4		
Total	68		

Marks
4

- The nickel(II) ion exists as the $[\text{Ni}(\text{OH}_2)_6]^{2+}$ complex ion in aqueous solution. Define the term complex.

What is the name of this complex ion?

Why is such a solution acidic?

Write a balanced equation for the corresponding reaction.


- You have completed a number of titrations during your laboratory work. What is the difference between the 'end point' and the 'equivalence point' in a titration?

4

How do you need to consider that distinction when you chose an indicator for a particular titration?

Marks
6

- Examine the following pressure/temperature phase diagram for a one component system.



Which phase exists in the fields labelled **A**, **B** and **C**?

A:	B:	C:
-----------	-----------	-----------

Explain your assignment of these phases.

--

What do the lines in the diagram represent?

--


What happens when you move across a line either by changing temperature or pressure?

--

For a compound with this phase diagram, would the solid be denser than the liquid or vice versa? Explain your answer.

--

- | |
|--------------|
| Marks |
| 4 |



RATE LAW	RATE CONSTANT
Answer:	Answer:

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

Marks
1

- Calculate the pH of a 0.020 M solution of $\text{Ba}(\text{OH})_2$.

pH =

3

- Calculate the pH of a 0.150 M solution of HNO_2 . The $\text{p}K_a$ of HNO_2 is 3.15.

pH =

2

- Calculate the pH of a solution that is 0.080 M in acetic acid and 0.160 M in sodium acetate. The $\text{p}K_a$ of acetic acid is 4.76.

pH =

Marks
4

- Hydrogen bonding is important for the physical properties of water and consequently the very existence of life on earth. What effect does the formation of hydrogen bonding have on the density of solid water (ice) compared to liquid water. Explain.

Predict the physical form of water under ambient conditions if no hydrogen bonds existed. Explain that prediction.

- BaSO_4 is used as a contrast agent for X-ray images of intestines. What is the solubility product constant, K_{sp} , for BaSO_4 , given that a maximum of 1.2×10^{-3} g dissolves in 500.0 mL of water.

Answer:






Ba^{2+} ions are toxic. Comment on the suitability of BaSO_4 as a contrast agent.

What advantage would there be in administering BaSO_4 as a slurry which also contains 0.5 M Na_2SO_4 ?

4


Marks
8

- Give the name of the starting material where indicated and the constitutional formula of the major organic product formed in each of the following reactions.






**Name:****Name:****Name:**

Marks
6

- Classify the starting materials for each of the following reactions as nucleophile and electrophile in the boxes provided and draw the structure of the product.



+




+

**THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.**

Marks
6

- Consider the following reaction sequence.







Give the reagents **C**, **D** and **F** and draw the structures of the major organic products, **A**, **B** and **E**, formed in these reactions.

A	D
B	E
C	F

Marks
5


- Show clearly the reagents you would use to carry out the following chemical conversion. Two steps are required. Give the structure of the intermediate compound.



How could you distinguish between the starting material and the product by ^{13}C NMR spectroscopy?

Marks
7

- Bupivacaine is the active molecule in some local anaesthetics. Of the two enantiomers, the one shown below (**X**) is the more effective.



What is the molecular formula of (**X**)?

Calculate the m/z value for the major peak you would expect to see for the molecular ion in the high resolution mass spectrum.

[Atomic masses: $^1\text{H} = 1.0078$; $^{12}\text{C} = 12.0000$; $^{16}\text{O} = 15.9949$; $^{14}\text{N} = 14.0031$]

Answer:

List the substituents attached to the stereogenic centre in descending order of priority according to the sequence rule.

highest priority

lowest priority


--	--	--	--

What is the absolute stereochemistry of (**X**)? Write (*R*) or (*S*).

Name the functional groups present in (**X**).

Marks
4

- Threonine (**Y**) is an amino acid. On the structure of (**Y**) below, identify all stereocentres in threonine with an asterisk (*).



How many possible stereoisomers of threonine are there?

Give the structures of the products obtained when threonine is treated with the following reagents.

1 M HCl

1 M NaOH

THE REMAINDER OF THIS PAGE IS FOR ROUGH WORKING ONLY.

CHEM1002 - FUNDAMENTALS OF CHEMISTRY 1B
DATA SHEET

Physical constants

Avogadro constant, $N_A = 6.022 \times 10^{23} \text{ mol}^{-1}$

Faraday constant, $F = 96485 \text{ C mol}^{-1}$

Planck constant, $h = 6.626 \times 10^{-34} \text{ J s}$

Speed of light in vacuum, $c = 2.998 \times 10^8 \text{ m s}^{-1}$

Rydberg constant, $E_R = 2.18 \times 10^{-18} \text{ J}$

Boltzmann constant, $k_B = 1.381 \times 10^{-23} \text{ J K}^{-1}$

Permittivity of a vacuum, $\epsilon_0 = 8.854 \times 10^{-12} \text{ C}^2 \text{ J}^{-1} \text{ m}^{-1}$

Gas constant, $R = 8.314 \text{ J K}^{-1} \text{ mol}^{-1}$
 $= 0.08206 \text{ L atm K}^{-1} \text{ mol}^{-1}$

Charge of electron, $e = 1.602 \times 10^{-19} \text{ C}$

Mass of electron, $m_e = 9.1094 \times 10^{-31} \text{ kg}$

Mass of proton, $m_p = 1.6726 \times 10^{-27} \text{ kg}$

Mass of neutron, $m_n = 1.6749 \times 10^{-27} \text{ kg}$

Properties of matter

Volume of 1 mole of ideal gas at 1 atm and 25 °C = 24.5 L

Volume of 1 mole of ideal gas at 1 atm and 0 °C = 22.4 L

Density of water at 298 K = 0.997 g cm⁻³

Conversion factors

1 atm = 760 mmHg = 101.3 kPa

1 Ci = $3.70 \times 10^{10} \text{ Bq}$

0 °C = 273 K

1 Hz = 1 s⁻¹

1 L = 10⁻³ m³

1 tonne = 10³ kg

1 Å = 10⁻¹⁰ m

1 W = 1 J s⁻¹

1 eV = $1.602 \times 10^{-19} \text{ J}$

Decimal fractions

Fraction	Prefix	Symbol
10 ⁻³	milli	m
10 ⁻⁶	micro	μ
10 ⁻⁹	nano	n
10 ⁻¹²	pico	p

Decimal multiples

Multiple	Prefix	Symbol
10 ³	kilo	k
10 ⁶	mega	M
10 ⁹	giga	G
10 ¹²	tera	T

CHEM1002 - FUNDAMENTALS OF CHEMISTRY 1B**Standard Reduction Potentials, E°**

Reaction	E° / V
$\text{S}_2\text{O}_8^{2-} + 2\text{e}^- \rightarrow 2\text{SO}_4^{2-}$	+2.01
$\text{Co}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Co}^{2+}(\text{aq})$	+1.82
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightarrow \text{Ce}^{3+}(\text{aq})$	+1.72
$\text{Au}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Au}(\text{s})$	+1.50
$\text{Cl}_2 + 2\text{e}^- \rightarrow 2\text{Cl}^-(\text{aq})$	+1.36
$\text{O}_2 + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightarrow 2\text{H}_2\text{O}$	+1.23
$\text{Br}_2 + 2\text{e}^- \rightarrow 2\text{Br}^-(\text{aq})$	+1.10
$\text{MnO}_2(\text{s}) + 4\text{H}^+(\text{aq}) + \text{e}^- \rightarrow \text{Mn}^{3+} + 2\text{H}_2\text{O}$	+0.96
$\text{Pd}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pd}(\text{s})$	+0.92
$\text{Ag}^+(\text{aq}) + \text{e}^- \rightarrow \text{Ag}(\text{s})$	+0.80
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightarrow \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{Cu}^+(\text{aq}) + \text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.53
$\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cu}(\text{s})$	+0.34
$\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}^{2+}(\text{aq})$	+0.15
$2\text{H}^+(\text{aq}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g})$	0 (by definition)
$\text{Fe}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.04
$\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Pb}(\text{s})$	-0.13
$\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Sn}(\text{s})$	-0.14
$\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ni}(\text{s})$	-0.24
$\text{Co}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Co}(\text{s})$	-0.28
$\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Fe}(\text{s})$	-0.44
$\text{Cr}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.74
$\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Zn}(\text{s})$	-0.76
$2\text{H}_2\text{O} + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$	-0.83
$\text{Cr}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Cr}(\text{s})$	-0.89
$\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightarrow \text{Al}(\text{s})$	-1.68
$\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Mg}(\text{s})$	-2.36
$\text{Na}^+(\text{aq}) + \text{e}^- \rightarrow \text{Na}(\text{s})$	-2.71
$\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ca}(\text{s})$	-2.87
$\text{Li}^+(\text{aq}) + \text{e}^- \rightarrow \text{Li}(\text{s})$	-3.04

CHEM1002 - FUNDAMENTALS OF CHEMISTRY 1B*Useful formulas*

Thermodynamics & Equilibrium $\Delta U = q + w = q - p\Delta V$ $\Delta_{\text{universe}}S = \Delta_{\text{sys}}S - \frac{\Delta_{\text{sys}}H}{T_{\text{sys}}}$ $\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ$ $\Delta G = \Delta G^\circ + RT \ln Q$ $\Delta G^\circ = -RT \ln K$ $K_p = K_c (RT)^{\Delta n}$	Electrochemistry $\Delta G^\circ = -nFE^\circ$ <i>Moles of e^- = It/F</i> $E = E^\circ - (RT/nF) \times 2.303 \log Q$ $= E^\circ - (RT/nF) \times \ln Q$ $E^\circ = (RT/nF) \times 2.303 \log K$ $= (RT/nF) \times \ln K$ $E = E^\circ - \frac{0.0592}{n} \log Q \text{ (at } 25^\circ\text{C)}$
Colligative properties $\pi = cRT$ $P_{\text{solution}} = X_{\text{solvent}} \times P^\circ_{\text{solvent}}$ $p = kc$ $\Delta T_f = K_f m$ $\Delta T_b = K_b m$	Quantum Chemistry $E = h\nu = hc/\lambda$ $\lambda = h/mv$ $4.5k_B T = hc/\lambda$ $E = -Z^2 E_R (1/n^2)$ $\Delta x \cdot \Delta(mv) \geq h/4\pi$ $q = 4\pi r^2 \times 5.67 \times 10^{-8} \times T^4$
Acids and Bases $pK_w = \text{pH} + \text{pOH} = 14.00$ $pK_w = \text{p}K_a + \text{p}K_b = 14.00$ $\text{pH} = \text{p}K_a + \log\{[A^-] / [HA]\}$	Gas Laws $PV = nRT$ $(P + n^2a/V^2)(V - nb) = nRT$
Radioactivity $t_{1/2} = \ln 2 / \lambda$ $A = \lambda N$ $\ln(N_0/N_t) = \lambda t$ $^{14}\text{C age} = 8033 \ln(A_0/A_t)$	Kinetics $t_{1/2} = \ln 2 / k$ $k = Ae^{-E_a/RT}$ $\ln[A] = \ln[A]_0 - kt$ $\ln \frac{k_2}{k_1} = \frac{E_a}{R} \left(\frac{1}{T_1} - \frac{1}{T_2} \right)$
Miscellaneous $A = -\log_{10} \frac{I}{I_0}$ $A = \epsilon cl$ $E = -A \frac{e^2}{4\pi\epsilon_0 r} N_A$	Mathematics If $ax^2 + bx + c = 0$, then $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $\ln x = 2.303 \log x$

PERIODIC TABLE OF THE ELEMENTS

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
1 HYDROGEN H 1.008																	2 HELIUM He 4.003
3 LITHIUM Li 6.941	4 BERYLLIUM Be 9.012											5 BORON B 10.81	6 CARBON C 12.01	7 NITROGEN N 14.01	8 OXYGEN O 16.00	9 FLUORINE F 19.00	10 NEON Ne 20.18
11 SODIUM Na 22.99	12 MAGNESIUM Mg 24.31											13 ALUMINIUM Al 26.98	14 SILICON Si 28.09	15 PHOSPHORUS P 30.97	16 SULFUR S 32.07	17 CHLORINE Cl 35.45	18 ARGON Ar 39.95
19 POTASSIUM K 39.10	20 CALCIUM Ca 40.08	21 SCANDIUM Sc 44.96	22 TITANIUM Ti 47.88	23 VANADIUM V 50.94	24 CHROMIUM Cr 52.00	25 MANGANESE Mn 54.94	26 IRON Fe 55.85	27 COBALT Co 58.93	28 NICKEL Ni 58.69	29 COPPER Cu 63.55	30 ZINC Zn 65.39	31 GALLIUM Ga 69.72	32 GERMANIUM Ge 72.59	33 ARSENIC As 74.92	34 SELENIUM Se 78.96	35 BROMINE Br 79.90	36 KRYPTON Kr 83.80
37 RUBIDIUM Rb 85.47	38 STRONTIUM Sr 87.62	39 YTTRIUM Y 88.91	40 ZIRCONIUM Zr 91.22	41 NIOBIUM Nb 92.91	42 MOLYBDENUM Mo 95.94	43 TECHNETIUM Tc [98.91]	44 RUTHENIUM Ru 101.07	45 RHODIUM Rh 102.91	46 PALLADIUM Pd 106.4	47 SILVER Ag 107.87	48 CADMIUM Cd 112.40	49 INDIUM In 114.82	50 TIN Sn 118.69	51 ANTIMONY Sb 121.75	52 TELLURIUM Te 127.60	53 IODINE I 126.90	54 XENON Xe 131.30
55 CAESIUM Cs 132.91	56 BARIUM Ba 137.34	57-71	72 HAFNIUM Hf 178.49	73 TANTALUM Ta 180.95	74 TUNGSTEN W 183.85	75 RHENIUM Re 186.2	76 OSMIUM Os 190.2	77 IRIDIUM Ir 192.22	78 PLATINUM Pt 195.09	79 GOLD Au 196.97	80 MERCURY Hg 200.59	81 THALLIUM Tl 204.37	82 LEAD Pb 207.2	83 BISMUTH Bi 208.98	84 POLONIUM Po [210.0]	85 ASTATINE At [210.0]	86 RADON Rn [222.0]
87 FRANCIUM Fr [223.0]	88 RADIUM Ra [226.0]	89-103	104 RUTHERFORDIUM Rf [261]	105 DUBNIUM Db [262]	106 SEABORGIUM Sg [266]	107 BOHRIUM Bh [262]	108 HASSIUM Hs [265]	109 MEITNERIUM Mt [266]	110 DARMSTADTIUM Ds [271]	111 ROENTGENIUM Rg [272]							

LANTHANOID
S

57 LANTHANUM La 138.91	58 CERIUM Ce 140.12	59 PRASEODYMIUM Pr 140.91	60 NEODYMIUM Nd 144.24	61 PROMETHIUM Pm [144.9]	62 SAMARIUM Sm 150.4	63 EUROPIUM Eu 151.96	64 GADOLINIUM Gd 157.25	65 TERBIUM Tb 158.93	66 DYSPROSIUM Dy 162.50	67 HOLMIUM Ho 164.93	68 ERBIUM Er 167.26	69 THULIUM Tm 168.93	70 YTTERBIUM Yb 173.04	71 LUTETIUM Lu 174.97
89 ACTINIUM Ac [227.0]	90 THORIUM Th 232.04	91 PROTACTINIUM Pa [231.0]	92 URANIUM U 238.03	93 NEPTUNIUM Np [237.0]	94 PLUTONIUM Pu [239.1]	95 AMERICIUM Am [243.1]	96 CURIUM Cm [247.1]	97 BERKELLIUM Bk [247.1]	98 CALIFORNIUM Cf [252.1]	99 EINSTEINIUM Es [252.1]	100 FERMIUM Fm [257.1]	101 MENDELEVIUM Md [256.1]	102 NOBELIUM No [259.1]	103 LAWRENCIUM Lr [260.1]

ACTINOIDS