

UNIVERSITY COLLEGE LONDON

University of London

EXAMINATION FOR INTERNAL STUDENTS

For The Following Qualifications:–

B.Sc. *M.Sc.*

Mathematics C348: Mathematics For General Relativity

COURSE CODE : MATHC348

UNIT VALUE : 0.50

DATE : 19–MAY–04

TIME : 14.30

TIME ALLOWED : 2 Hours

All questions may be attempted but only marks obtained on the best **four** solutions will count.

The use of an electronic calculator is **not** permitted in this examination.

Geodesic Equation ($U^a = dX^a/d\tau$)

$$\frac{dU^a}{d\tau} + \Gamma^a_{bc} U^b U^c = 0; \quad (1)$$

$$\frac{dp_a}{d\tau} = \frac{m}{2} (\partial_a g_{bc}) U^b U^c; \quad (2)$$

$$\Gamma^a_{bc} = \frac{1}{2} g^{ad} (\partial_b g_{cd} + \partial_c g_{bd} - \partial_d g_{bc}). \quad (3)$$

Geodesics parameterized by λ

$$\frac{d^2 X^a}{d\lambda^2} + \Gamma^a_{bc} \frac{dX^b}{d\lambda} \frac{dX^c}{d\lambda} = 0. \quad (4)$$

Schwarzschild metric line element

$$d\tau^2 = \left(1 - \frac{r_s}{r}\right) dt^2 - \left(1 - \frac{r_s}{r}\right)^{-1} dr^2 - r^2 (d\theta^2 + \sin^2 \theta d\phi^2); \quad r_s = 2GM.$$

Faraday tensor

$$F_{ab} = \begin{pmatrix} 0 & -E_x & -E_y & -E_z \\ E_x & 0 & B_z & -B_y \\ E_y & -B_z & 0 & B_x \\ E_z & B_y & -B_x & 0 \end{pmatrix}; \quad F^{ab} = \begin{pmatrix} 0 & E_x & E_y & E_z \\ -E_x & 0 & B_z & -B_y \\ -E_y & -B_z & 0 & B_x \\ -E_z & B_y & -B_x & 0 \end{pmatrix}.$$

Dual Faraday tensor

$$*F^{ab} \equiv \frac{1}{2} \epsilon^{abcd} F_{cd} = \begin{pmatrix} 0 & +B_x & +B_y & +B_z \\ -B_x & 0 & -E_z & E_y \\ -B_y & E_z & 0 & -E_x \\ -B_z & -E_y & E_x & 0 \end{pmatrix}.$$

Maxwell Source Equations

$$\partial_b F^{ab} = j_e^a.$$

Internal Maxwell Equations

$$\partial_a F_{bc} + \partial_b F_{ca} + \partial_c F_{ab} = 0 \quad \text{or} \quad \partial_b *F^{ab} = 0.$$

1. (a) How does a second rank mixed tensor Q^a_b transform under a coordinate transformation between unprimed coordinates x^a and primed coordinates x'^a ?
- (b) The trace of a second rank mixed tensor T is defined by

$$\text{trace}(T) = T^a_a.$$

Show using the transformation laws, that $\text{trace}(T)$ is a scalar, i.e. that it takes the same value in all coordinate frames.

- (c) Given a metric tensor g_{ab} , what is the definition of the tensor g^{cd} ? What is the tensor $g_{ab}g^{bc}$?
- (d) Suppose the metric tensor g_{ab} has both a symmetric part $g_{Sab} = g_{Sba}$ and an antisymmetric part, $g_{Aab} = -g_{Aba}$,

$$g_{ab} = g_{Sab} + g_{Aab}.$$

Show explicitly that the line element $d\tau^2$ is independent of g_{Aab} .

- (e) Consider a two-dimensional manifold M with coordinates x^1 and x^2 . Suppose the metric is

$$g_{ab} = \begin{pmatrix} 3 & 0 \\ 0 & 2 \end{pmatrix}.$$

Also let A^a be a vector and S^a_{bc} a tensor with values

$$\begin{aligned} A^1 &= -1, & A^2 &= 2; \\ S^1_{11} &= 1, & S^1_{12} &= 0, & S^1_{21} &= 2, & S^1_{22} &= 3, \\ S^2_{11} &= 0, & S^2_{12} &= 1, & S^2_{21} &= 2, & S^2_{22} &= 2. \end{aligned}$$

Find the following tensors:

- (i) A_b ;
- (ii) $X_c = S^a_{ac}$;
- (iii) $B^a_c = X^a X_b$.
- (f) Using index notation prove the vector identity

$$\nabla \cdot (A \times B) = B \cdot \nabla \times A - A \cdot \nabla \times B.$$

- (g) Determine which of the following equations is a valid tensor equation. Describe the errors in the other equations.

- (i) $A_{ab} = B^{ab}F^c_c$;
- (ii) $W^a_{bc} = D_c E_b F^a$;
- (iii) $f = J_b K_a L^a$.

2. (a) What is the definition of the Lorentz group? Show that all Lorentz transformations have determinant ± 1 . Give an example of a Lorentz transformation with determinant -1 .

- (b) Suppose inertial reference frame B moves at a velocity V in the x direction with respect to inertial frame A.

Derive the Lorentz transformation between A and B, using the principles of special relativity. You may assume that the transformation is linear, so that

$$\begin{pmatrix} t \\ x \end{pmatrix}_B = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} t \\ x \end{pmatrix}_A$$

for some numbers a, b, c, d . You may also assume that $y_A = y_B, z_A = z_B$.

- (c) The Faraday Tensor F_{ab} can be written in terms of the 4-vector potential ϕ_a as $F_{ab} = \partial_b \phi_a - \partial_a \phi_b$. The Maxwell Source Equations are $\partial_b F^{ab} = j^a$ where j^a is the current density.

- (i) Show that $\partial_a j^a = 0$. Describe the physical meaning of this equation.
(ii) Write the Maxwell Source Equations in terms of ϕ^a instead of F^{ab} . What do the equations reduce to if we assume Lorentz gauge, $\partial_a \phi^a = 0$?

3. The covariant derivative of a covariant second rank tensor M_{cd} is

$$(\nabla_b M)_{cd} = \partial_b M_{cd} - \Gamma^a_{bc} M_{ad} - \Gamma^a_{bd} M_{ca}.$$

Christoffel symbols are assumed to be symmetric in their last two indices:

$$\Gamma^a_{cb} = \Gamma^a_{bc}.$$

- (a) Given that the covariant derivative of the metric is always zero, i.e. $(\nabla_b g)_{cd} = 0$ for any choice of b, c , and d , derive the equation

$$\Gamma^a_{bc} = \frac{1}{2} g^{ad} (\partial_b g_{cd} + \partial_c g_{bd} - \partial_d g_{bc}).$$

- (b) Consider Euclidean 3-space \mathbb{E}^3 with cylindrical coordinates (ρ, ϕ, z) . A conical surface within this space can be defined by the equation

$$z = a\rho$$

with a constant. Let the coordinates on the cone be $(C^1, C^2) = (\rho, \phi)$. Show that the metric for the cone in these coordinates is

$$g_{ab} = \begin{pmatrix} 1 + a^2 & 0 \\ 0 & \rho^2 \end{pmatrix}.$$

- (c) Calculate the eight Christoffel symbols Γ^a_{bc} for this metric.
(d) Consider a geodesic on this cone, parameterized by arclength s . Find expressions for $d^2\phi/ds^2$ and $d^2\rho/ds^2$ along the geodesic.

4. The Newtonian orbit equation for a planet at radius r as a function of angle ϕ is

$$\frac{d^2u}{d\phi^2} + u = \frac{GM}{h^2},$$

where $u = 1/r$ and $h = \text{angular momentum/mass}$. Geodesics in the Schwarzschild metric satisfy the equation

$$\frac{d^2u}{d\phi^2} + u = \frac{GM}{h^2} + (3GM)u^2.$$

- (a) Consider a planet of mass m moving in a geodesic around the sun (mass M) whose gravitational field is described by the Schwarzschild metric. The planet moves in the ecliptic plane $\theta = \pi/2$. What is the Schwarzschild metric in this plane? Obtain expressions for $dt/d\tau$ and $d\phi/d\tau$ in terms of two conserved quantities energy/mass k and angular momentum/mass h .
- (b) Show that

$$u(\phi) = u_0(1 + \epsilon \sin \phi)$$

is a solution of the Newtonian orbit equation. Find u_0 in terms of r_s and h . At what angle ϕ is perihelion for this solution? What kind of orbit has $\epsilon = 0$?

- (c) Let

$$u(\phi) = u_0(1 + \epsilon \sin \phi + y(\phi))$$

where $y(\phi) \ll 1$. Consider an orbit with $\epsilon \ll 1$. Find an approximate differential equation for $y(\phi)$ good to first order in ϵ and y .

- (d) What are the complementary functions for the differential equation derived in (c)? What are the particular integrals? Find the solution given conditions at perihelion $y(\pi/2) = 0$, $y'(\pi/2) = 0$.
- (e) The solution for $y(\phi)$ includes the precession term

$$(3GM\epsilon u_0) \left(\frac{\pi}{2} - \phi \right) \cos \phi,$$

plus periodic terms. (Hint: these periodic terms can be neglected). Let the next perihelion be at $\phi = 5\pi/2 + \delta\phi$ where $\delta\phi \ll 1$. Give an approximate expression for the precession angle $\delta\phi$.

5. A spaceship accelerates away from Earth in the x direction. The astronauts feel a constant acceleration equal to g , the acceleration at the Earth's surface. For this question, ignore the y and z coordinates. Proper time on the ship is given by τ . The speed of the ship in the Earth's frame is $V(\tau)$. Let the transformation from ship coordinates S to Earth coordinates E be

$$\frac{\partial E^a}{\partial S^b} = \begin{pmatrix} \partial t_E / \partial t_S & \partial t_E / \partial x_S \\ \partial x_E / \partial t_S & \partial x_E / \partial x_S \end{pmatrix} = \begin{pmatrix} \gamma(\tau) & \gamma(\tau)V(\tau) \\ \gamma(\tau)V(\tau) & \gamma(\tau) \end{pmatrix}$$

where $\gamma(\tau) = (1 - V^2(\tau))^{-1/2}$. The velocity 4-vector in the ship's frame is

$$u_S^a(\tau) = \begin{pmatrix} 1 \\ 0 \end{pmatrix}.$$

The covariant form of Newton's second law of motion for an object of mass m subject to a force F^a is

$$m \frac{Du^a}{D\tau} = m \left(\frac{du^a}{d\tau} + \Gamma^a_{bc} u^b u^c \right) = F^a.$$

- The acceleration at the Earth's surface is approximately $g = 10 \text{ m s}^{-2}$. Derive an approximate expression for this quantity in relativistic units, i.e. in units of year^{-1} .
- Let $\phi(\tau)$ be defined by $V(\tau) = \tanh \phi(\tau)$. Write the transformation matrix in terms of $\phi(\tau)$. What is $u_E^a(\tau)$? What is $du_E^a(\tau)/d\tau$?
- In the ship's frame, $F_S^a = \begin{pmatrix} 0 \\ mg \end{pmatrix}$. What is $Du_S^a/D\tau$ in the ship's frame? What is Γ^1_{00} ?
- What is $Du_E^a/D\tau$ in the Earth frame? If we suppose that the Earth frame is a Locally Inertial Frame, what is Γ^a_{Ebc} ? Hence find a second expression for $du_E^a/d\tau$ in the Earth's frame.
- Using the results above, find the function $\phi(\tau)$. Hence find the function $t_E(\tau)$.