

UNIVERSITY OF KWAZULU-NATAL

SUPPLEMENTARY EXAMINATION: 12th January 2009

Subject, Course and Code: Physics Foundation PHYS099 and PHYS199

DURATION: 3 HOURS

TOTAL MARKS:180

Seat Number: _____

Student Number: _____

INTERNAL EXAMINERS: Ms W. Dlamini, Mrs S. Halstead, Ms A. Marais, Mr R. Webber and Mr E. Zhandire

EXTERNAL EXAMINER: Dr V. Couling

ANSWER ALL QUESTIONS ON THIS PAPER

IN THEIR OWN INTERESTS STUDENTS ARE REQUESTED TO WRITE LEGIBLY.

THIS PAPER CONSISTS OF 12 PAGES. PLEASE SEE THAT YOU HAVE THEM ALL.

Properties of Matter

Question 1 (14 marks)

Give your answers to the following questions with the correct number of significant figures or decimal places.

(a) Calculate the mass of 98 identical pieces of copper if the mass of each piece is 0.0508 kg. (2)

(b) $1.05 \text{ m} + 1005 \text{ cm} - 1050 \text{ mm}$ (2)


(c) $120.5 \text{ km/h} = \underline{\hspace{4cm}} \text{ m/s}$ (2)

(d) $(11.06 \pm 0.05) \text{ cm} - (90.5 \pm 0.3) \text{ mm}$ (3)

(e) A room is 16.40 m long by 4.5 m wide by 3.26 m high. The density of air is 1.29 kg/m^3 . Find the mass of the air contained in the room. (5)

Kinematics**Question 2 (9 marks)**


The following velocity-time graph represents the movement of a car that was **initially travelling towards the East**.



- (a) At which stage(s) did the car have a constant velocity? (1)
- (b) Use the graph to calculate the acceleration from E to F. (3)
- (c) Use the graph to determine the displacement of the car between A and C. (3)
- (d) Describe the motion of the car from B to C. (2)

Question 3 (13 marks)

A man throws a ball upwards at an angle of 30.0° to the horizontal at a speed of 5.00 ms^{-1} from the top of a building 6.00 m above the ground, as shown in the diagram below. Taking upwards as positive and the magnitude of $g = 9.80 \text{ ms}^{-2}$ calculate, to the correct number of significant figures:




(a) the magnitude and direction of the ball's vertical velocity as it reaches the ground, (5)

(b) the time taken for the ball to reach the ground, and (4)

(c) how far from the base of the building, in the horizontal direction, the ball lands. (4)

Question 4 (11 marks)

A towel is held on an inextensible washing line, as shown alongside. The tensions T_1 and T_2 are 12 N and 13.5 N respectively. Take the magnitude of $g = 9.80 \text{ ms}^{-2}$.



(a) Calculate the mass of the towel. (6)

(b) The cord of the washing line has a maximum tensile strength of 25N. Suppose the mass of the towel doubles when it is wet, and all angles remain the same. Will the cord be able to support the weight of the wet towel if it is hung in the same position? (The maximum tensile strength of the cord is the maximum amount of tension the cord can support without breaking.) (5)

Question 5 Astronomy

Next to the term in column A, write the letter of the term in column B, which best describes it.

Column A

Planets _____
 Proxima Centauri _____
 Sun _____
 Moon _____
 Milky Way _____

Column B


a) A satellite
 b) Galaxy
 c) Move around a sun
 d) Star
 e) Produces light

Electrostatics Take the electrostatic constant $k = 9.00 \times 10^9 \text{ NmC}^{-2}$

Question 6 (30 marks)

- (a) Two charged objects have a repulsive force of 0.0980 N. If the distance between the two charges is doubled, then what is the new force? (3)

- (b) Three charges Q_1 , Q_2 and Q_3 with positions shown in the diagram exert electrostatic forces on each other.




- (i) Determine the magnitude and direction of the resultant electrostatic force on Q_2 . (11)

- (ii) Determine the magnitude and direction of the electric field at the point where Q_2 is situated. (4)

(c) Two parallel charged plates, X and Y, are placed as shown in the diagram. Consider only electrostatic effects.

- (i) If a positive charge is allowed to move freely, horizontally from Y to X, what energy changes take place? (2)



- (ii) If a positive charge is moved vertically from L to M, what energy changes take place? (1)


(d) A charge Q of $4.0 \mu\text{C}$ exerts a force of $3.0 \times 10^{-2} \text{ N}$ to the right, on a charge q of 2.0 nC . Calculate: (Q) (q)

- (i) The electric field at q (4)

- (ii) The distance between the charges (5)

Electric circuits**Question 7 (21 marks)**


The following circuit has **identical resistors each of resistance R** . The battery and connecting leads have no resistance. There is a current of 400 mA as shown in the diagram, and the voltage across resistor D is 1.50 V.



- (a) What is the current through:
- (i) Resistor C? (1)
 - (ii) Resistor D? (3)
 - (iii) Resistor A? (2)
- (b) Calculate amount of charge that flows through resistor E in 20 s. (3)
- (c) Calculate
- (i) The voltage across resistor C (3)
 - (ii) The voltage across resistor A. (3)
 - (iii) The battery voltage (3)
 - (iv) The resistance R of one resistor. (3)

Question 8 (6 marks)

Hlonophile has R1.05 credit on her pre-paid electricity meter. Electricity cost 50c per unit of kilowatt-hour. She gets home, cooks supper before dark using a 2000W stove for 50 minutes, then studies, using a 100 W light bulb. How long will she be able to study? Show calculations to support your answer. (6)


Question 9 (10 marks)

The diagram above shows an electrical circuit which consists of an unknown resistor R_x connected in series with a $25.0\ \Omega$ resistor and a battery of emf of $12.0\ \text{V}$ and internal resistance $0.320\ \Omega$. An ammeter connected in the circuit reads $200\ \text{mA}$.

(a) A voltmeter is connected across the battery when the circuit is closed. What is its reading? (3)

(b) Calculate the value of the unknown resistor R_x . (4)

(c) Calculate the rate at which energy is used by the $25.0\ \Omega$ resistor. (3)

Question 10 (8 marks)

In the circuit shown above the unknown resistances R_x and R_y are equal. Use the nodes and loops marked to find:

- (a) Current, I_x (2)
- (b) Resistance, R_x (3)
- (c) Battery voltage, V_x (3)

Newton's Laws**Question 11 (7 marks)**


An astronaut lands on a spherical (round) planet, X, of radius 150 km. She finds that a 2.0 kg mass released from rest at a height of 1.0 m takes 1.5 s to reach the surface.

Take $G = 6.67 \times 10^{-11} \text{ Nm}^2\text{kg}^{-2}$. Calculate:

- (a) the gravitational acceleration of the object on planet X, and (3)
- (b) the mass of planet X. (4)


Question 12 (8 marks)

On the diagram alongside, draw in and label all the forces acting on the car as it is being pushed up the hill at a constant velocity. Show approximate magnitudes. (8)

**Question 13 (9 marks)**

A box is being dragged along a rough concrete floor, as shown in the diagram. The mass of the box is 50 kg and the force exerted by the rope is 200 N, while the force of friction is 25 N. Take “up” and “left” to be positive as shown in the diagram alongside. Determine:

(a) the normal force acting on the box, and (4)



(b) the acceleration of the box. (5)

Question 14 (8 marks)

Determine the braking force needed to bring a small truck of mass 2000 kg travelling **west** at 120 km/h to a stop in 100 m. (8)

Optics


Question 15 (21 marks)

- (a) The diagram below shows light entering a glass block which is placed above some water. Use the table of refractive indices to answer these questions. Please use a ruler, and show angles that are approximately correct, you do not need to calculate them.


material	refractive index
air	1.00
ice	1.31
water	1.33
perspex	1.42
glass	1.58

On the diagram


- (i) Mark in the angle of incidence. (1)
- (ii) Show the path of the light right through into the water. (3)



- (b) The surfaces of a plane mirror and a rectangular block of Perspex are at right angles to each other as shown in the diagram below. The angle of refraction of a light ray is measured as 21.5° in the Perspex. Calculate the angle of incidence of the light ray, θ , at the surface of the mirror. Explain your reasoning at each stage. (6)



- (c) A periscope makes use of total internal reflection. It uses a 45° triangular glass prism such as shown in the diagram below. Use the data from the table on the previous page to help answer these questions.



- (i) **On the diagram above** label clearly
- (A) At least one surface where no bending of the light ray takes place. (1)
- (B) At least one place where total internal reflection will take place. (2)
- (ii) Calculate the critical angle for glass. (3)
- (iii) If the prism in the diagram were made of ice, would the periscope work? Explain by means of a calculation. (5)

TOTAL MARKS 180